

# FGFR1c Human Recombinant Antibody (MM-0307)



Catalog #: CR - 008

Size: 100 µg

## Product Information


**Name:** FGFR1c Human Recombinant Antibody  
**Antigen:** Human FGFR1c (ECD: Arg22 – Glu374)  
**Isotype:** Human IgG1; custom orders also available as mouse IgG1, single chain IgG1 & single chain human-mouse IgG2a  
**Reactivity:** Human, mouse  
**Clonality:** Monoclonal  
**Source:** Human (Expi293 cell line by Life Technologies)  
**Applications:** ELISA and Flow Cytometry

**Concentration:** Lyophilized protein A-purified Antibody in DPBS pH 7.4; no preservatives added

**Handling:** Lyophilized antibody can be kept at 4°C for 3 months. To avoid freeze-thaw cycles, reconstituted antibody should be resuspended in H<sub>2</sub>O and aliquoted before freezing for short term (-20°C) or for long term storage (-80°C). For maximum recovery of products, centrifuge the vial prior to removing the cap. Further dilutions can be made in assay buffer. A final stock concentration of 1 µg/µl is recommended.

## Representative data


### Capture ELISA



#### Capture ELISA of FGFR1c:

1 µg/mL of human or mouse FGFR proteins or control proteins (Fc and BSA) were adsorbed onto maxisorp plates, blocked with PBS + 0.5% BSA (PB), and washed 3 times with PBS + 0.05% Tween-20 (PT); 1 µg/ml anti-FGFR1c Fab antibody in PBS for 30 min, shaking RT, washed 6 times with PT. Anti-human kappa-HRP (1:5000 in PB) was incubated for 30 min, shaking RT, washed 6 times with PT. Colorimetric HRP reagents allow for absorbance readings at 450 nm.

### Flow Cytometry



#### Flow Cytometry of FGFR1c:

H1581 cells (were collected using a mild EDTA solution and washed in PBS. 5x10<sup>5</sup> cells were incubated with 0.1 µg of anti-FGFR1c hIgG1, antibody in 100 µL PBS + 0.1% BSA for 30 min on ice, then washed twice with cold PBS + 0.1% BSA. Sample was then incubated with Goat anti-Human Fab-488 (Jackson ImmunoResearch, diluted with 1:1000 PBS + 0.1% BSA) for 15-30 min, washed twice with cold PBS + 0.1% BSA and fixed with 2% paraformaldehyde and filtered prior to reading on a FACS machine.

*\*Antibody staining is not recommended after fixing cells\**

*Optimal dilution of the antibody to be determined by the user*

## Limitations and Liability

For research use only not for human use; Manufactured by CCAB under a license for use of Expi293 cells (Life Technologies). CCAB is not responsible under any circumstances for any lost profit, loss of business, or any form of indirect, special, incidental, consequential, punitive, or treble damages, however caused, arising out of or relating to use of this product. This product is not for resale. Any modification of the antibody for commercial purposes is prohibited.